Specific health screenings.

Targeted Cancer Screening



We'll focus on spotting signs of some of the most common cancers and support with any next steps, without the need for a GP referral.





Recommended for:

■ Those aged 18 and over*

*HPV/Smear tests are not clinically recommended for those under 25. Under 25's can discuss their concerns with a doctor who will advise on best next steps if there are medical concerns about this.



Times

In-centre consultation with a GP - up to 30 minutes

Testicular Health

Advice on the signs and symptoms of testicular cancer and examination if needed.

- Review of medical history relating to testicular cancer
- Testicular examination¹

Breast Health

Get a picture of your current breast health and discuss any potential concerns with a GP.

- Full consultation including medical history review related to breast cancer
- Breast examination¹ if clinically indicated
- Onward referral during the appointment or upon receipt of any test results, if needed.

Prostate Health

Advice on the signs and symptoms of prostate cancer, plus a PSA blood test and examination if needed.

- Review of medical history relating to prostate cancer
- PSA Prostate cancer screening blood test. Aged 50+1
- Prostate examination¹
- Onward referral during the appointment or once the PSA results are in, if needed.

Cervical Health

Get a picture of your current cervical health and discuss any potential concerns with a GP.

- Full consultation including medical history review related to cervical cancer
- Cervical examination¹ if clinically indicated
- Test for human papilloma virus (HPV) for those aged 25 and over²
- Onward referral during the appointment or upon receipt of any test results, if needed.

Add ons

- Mammography: Aged 40+
- Bowel cancer screening test (stool) 45+
- Colon CT (Computerized Tomography) 45+

¹ Based on the results of the health questionnaire, depending on risk factors and medical history. Some tests may only be provided if considered appropriate by the doctor.

² Smear will be carried out if HPV virus is present, in line with NHS guidelines for best practice